

PROHIBIT[®]

(Levamisole Hydrochloride)

Soluble Drench Powder

Prohibit Soluble Powder is a broad spectrum anthelmintic effective against stomach, intestinal and lungworms in beef cattle and sheep.

- Available in 52 gram pouch and 605 gram bottle
- Broad coverage of nine major types of worms
- Short withdrawal time
- Low volume dosage: cattle - 2 mL/100 lbs.; sheep - 1 mL/50 lbs.
- Cost-effective and well-targeted
- For oral use in cattle and sheep



(30 x 52 gram pouch bucket)

Directions and Dosages

Cattle and sheep de-wormer. For oral use.

CATTLE STANDARD DRENCH SOLUTION - Place the contents of each pouch in a 1 quart (32 fl. oz.) container, fill with water; swirl until dissolved. Administer as a single drench dose according to the table on reverse side.

SHEEP STANDARD DRENCH SOLUTION - Place the contents of each pouch in a one gallon (128 fl. oz.) container, fill with water; swirl until dissolved. Administer as a single drench dose according to the table on reverse side.

Read full label and directions before use.



PROHIBIT[®] Soluble Drench Powder

Levamisole Hydrochloride Anthelmintic

Each 52 g pouch contains 46.8 g of levamisole hydrochloride activity.

ANADA 200-386, Approved by FDA

Each 605 g bottle contains 544.5 g of levamisole hydrochloride activity.

ANADA 200-225, Approved by FDA

Cattle and Sheep Dewormer. For Oral Use.

Administer as a standard drench with standard drench syringe or administer as a concentrated drench solution with an automatic drenching syringe.

Mixes as a clear solution.

INDICATIONS:

Prohibit[®] (Levamisole Hydrochloride) is a broad-spectrum anthelmintic and is effective against the following nematode infections in cattle and sheep:

SHEEP: STOMACH WORMS: (*Haemonchus contortus*, *Trichostrongylus axei*, *Teladorsagia circumcincta*), INTESTINAL WORMS: (*Trichostrongylus colubrifomis*, *Cooperia curticei*, *Nematodirus spathiger*, *Bunostomum trigonocephalum*, *Oesophagostomum columbianum*, *Chabertia ovina*), LUNGWORMS: (*Dictyocaulus filaria*).

CATTLE: STOMACH WORMS: (*Haemonchus placei*, *Trichostrongylus axei*, *Ostertagia ostertagi*), INTESTINAL WORMS: (*Trichostrongylus longispicularis*, *Cooperia oncophora*, *Cooperia punctata*, *Nematodirus spathiger*, *Bunostomum phlebotomum*, *Oesophagostomum radiatum*), LUNGWORMS: (*Dictyocaulus viviparus*).

DOSAGE AND ADMINISTRATION

CATTLE-STANDARD DRENCH SOLUTION- Place the contents of each pouch in a 1 quart (32 fl. oz.) container, fill with water; swirl until dissolved. Administer as a single drench dose according to the following table:

Weight	Drench Dosage	Each Pouch Will Treat
200 lb	1/2 fl oz	64 head
400 lb	1 fl oz	32 head
600 lb	1 1/2 fl oz	21 head
800 lb	2 fl oz	16 head

CATTLE-CONCENTRATED DRENCH SOLUTION- For use with 20 mL Automatic Syringe. Place the contents of each pouch in a standard household measuring container and add water to the 8 3/4 fl oz level. Swirl until dissolved. Give 2 mL (milliliter) per 100 lb body weight. Refer to the table above for the number of cattle each pouch will treat.

SHEEP-STANDARD DRENCH SOLUTION- Place the contents of each pouch in a 1 gallon (128 fl oz) container, fill with water; swirl until dissolved. Administer as a single drench dose according to the following table:

Weight	Drench Dosage	Each Pouch Will Treat
50 lb	1/2 fl oz	256 head
100 lb	1 fl oz	128 head
150 lb	1 1/2 fl oz	84 head
200 lb	2 fl oz	64 head

and Goats

SHEEP-CONCENTRATED DRENCH SOLUTION- For use with 20 mL automatic Syringe. Place the contents of each pouch in a standard household measuring container and add water to the 17 1/2 fl. oz. level. Swirl until dissolved. Give 2 mL per 50 lb. body weight. Refer to the table above for the number of sheep each pouch will treat.

GALLON BOTTLE - When you are ready to de-worm your cattle or sheep, add water to the powder in this bottle up to the 3 liter mark. Swirl to mix thoroughly before using. If any solution is left over, it may be stored for up to 3 months in this tightly capped bottle, shake well before using.

Administer as a single drench dose as follows:

Cattle -- 2 mL per 100 lb. body weight

Weight	Drench Dosage	Each Bottle Will Treat
100 lb	2 mL	1,500 head
300 lb	6 mL	500 head
500 lb	10 mL	300 head
700 lb	14 mL	214 head
1,000 lb.	20 mL	150 head

Sheep -- 1 mL per 50 lb. body weight and Goats

Weight	Drench Dosage	Each Bottle Will Treat
100 lb	2 mL	1,500 head
300 lb	6 mL	500 head
500 lb	10 mL	300 head
700 lb	14 mL	214 head
1,000 lb.	20 mL	150 head

NOTE: Careful weight estimates are essential for proper performance of this product. Prepare solutions as needed. However, excess solutions may be stored in clean closed containers up to 90 days without loss of anthelmintic activity.

Cattle and sheep maintained under conditions of constant helminth exposure may require retreatment within 2 to 4 weeks after the first treatment.

WARNING: Do not administer to cattle within 48 hours of slaughter for food. Do not administer to sheep within 72 hours of slaughter for food. To prevent residues in milk, do not administer to dairy animals of breeding age.

CAUTION: Muzzle foam may be observed. However, this reaction will disappear within a few hours. If this condition persists, a veterinarian should be consulted. Follow recommended dosage carefully. Consult veterinarian before using in severely debilitated animals.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Store between 15° and 30°C (59° and 86°F).

WARNING: Keep out of reach of children.

For Animal Use Only

® Registered trademark of Agri Laboratories, Ltd.

Manufactured for Agri Laboratories, Ltd., St. Joseph, MO 64503



P.O. Box 3103 St. Joseph, MO 64503

800-542-8916 www.AgriLabs.com

©2012 Agri Laboratories Ltd.

ANAD-0048 Rev 0213