

# Safe-Guard® Dewormer for Goats

## Intervet

Suspension 10%  
(100 mg/mL)

### Indications:

**Goats** - 2.3 mg/lb (5mg/kg) body weight for the removal and control of Stomach worms (adults): *Haemonchus contortus* and *Teladorsagia circumcincta*.


### Directions:

Determine the proper dose according to estimated body weight. Administer orally. The recommended dose of 2.3 mg/lb (5 mg/kg) is achieved when 2.3 mL of the drug are given for each 100 lb body weight.

<u>Body Weight</u>	<u>Amount</u>	<u>Double Amount</u>	<u>1 mL per 20 lbs</u>
25 lb	0.6 mL	1.2 mL	1.25 mL
50 lb	1.2 mL	2.4 mL	2.50 mL
75 lb	1.7 mL	3.4 mL	3.75 mL
100 lb	2.3 mL	4.6 mL	5.00 mL
125 lb	2.9 mL	5.8 mL	6.25 mL
150 lb	3.4 mL	6.8 mL	7.50 mL
175 lb	4.0 mL	8.0 mL	8.75 mL

Under conditions of continued exposure to parasites, retreatment may be needed after 4 to 6 weeks.

Store at or below 25°C (77°F). Protect from freezing. Shake well before use.



**RESIDUE WARNINGS: Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats.**

### Caution:

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Keep this and all medication out of the reach of children.

Intervet, Inc.  
405 State St., P.O. Box 318, Millsboro, DE 19966  
1-800-441-8272  
[www.intervet-usa.com](http://www.intervet-usa.com)

# Safe-Guard® Dewormer for Goats